



Nanosystems Initiative Munich



PRESS RELEASE

Energy saving chaperone Hsp90

Large conformational changes without ATP consumption

A special group of proteins, the so-called chaperones, helps other proteins to obtain their correct conformation. Until now scientists supposed that the chaperone Hsp90 needs energy from hydrolyzing ATP for its large conformational changes. Now a research team from the Nanosystems Initiative Munich (NIM) could prove that Hsp90 utilizes thermal fluctuations as the driving force for its conformational changes. The renowned journal PNAS reports on their findings.

ATP is the major energy source for most organisms and ATPases are the machines, which utilize this fuel, for example to move muscles or cargo in our body. The very abundant chaperone protein Hsp90 has such an ATPase in each of its two monomers. During the last years experiments had suggested that the movement and conformational changes of ATPase proteins are in general strictly linked to ATP binding and hydrolysis (i.e. fuel consumption).

To probe this theory Thorsten Hugel, Biophysics Professor at the Technische Universität München (TUM) and member of the Nanosystems Initiative Munich (NIM), and his team designed a particular three color single-molecule FRET (Förster resonance energy transfer) assay with alternating laser excitation (ALEX) for simultaneous observation of ATP binding and conformational changes. Unexpectedly the experiments revealed that binding and hydrolysis of ATP is not correlated with the large conformational changes of Hsp90. Hsp90 is instead a highly flexible machinery driven by thermal fluctuations.

"Thermal fluctuations are random changes in the structure of the protein – they can be thought of as collisions with water molecules in the environment, which move rather violently at the temperatures in a living organism," says Thorsten Hugel. "Using these clashes to switch back and forth between different conformations, saves Hsp90 valuable ATP." But then what is the task of ATPase in the Hsp90 chaperone? The scientists suspect that co-chaperones and substrate proteins alter the system so that ATP binding or hydrolysis can take a crucial task.

With the newly developed experimental setup, it is now possible to investigate the very complex system in greater detail to resolve this important question. The Munich biophysicists thereby offer a new perspective on the energy conversion in molecular machines.

Nanosystems Initiative Munich (NIM)
Dr. Birgit Gebauer (Presse- und Öffentlichkeitsarbeit)
Schellingstraße 4, D-80799 München
Tel.: +49 (89) 2180 5091 Fax: +49 (89) 2180 5649
Mail: birgit.gebauer@lmu.de

The work was supported by grants from the DFG (Hu997/9-1, SFB 863) and the Cluster of Excellence Nanosystems Initiative Munich (NIM) and the NanoBio-Technology program of the Elite Network of Bavaria.

Publication:

Heat shock protein 90's mechano-chemical cycle is dominated by thermal fluctuations.
Christoph Ratzke, Felix Berkemeier, and Thorsten Hugel (PNAS, January 3, 2012 vol. 109, no. 1, 161–166).

Contact:

Prof. Dr. Thorsten Hugel
Technische Universität München (TUM)
Department Biophysics and Institute for Medical Engineering
85748 Garching/Munich


Tel: +49 (0)89 / 289 16781

E-Mail: thugel@mytum.de

Website: <http://www.bio.ph.tum.de/index.php?id=6>

Picture:

ATP binds to the closed conformation of Hsp90



Nanosystems Initiative Munich (NIM)
Dr. Birgit Gebauer (Presse- und Öffentlichkeitsarbeit)
Schellingstraße 4, D-80799 München
Tel.: +49 (89) 2180 5091 Fax: +49 (89) 2180 5649
Mail: birgit.gebauer@lmu.de